arXiv:1303.2521v1 [math.DS] 11 Mar 2013

DYNAMICS OF ITERATED FUNCTION SYSTEMS ON THE CIRCLE CLOSE TO ROTATIONS

PABLO G. BARRIENTOS AND ARTEM RAIBEKAS

ABSTRACT. We study the dynamics of iteration function systems generated by a pair of circle diffeomorphisms close to rotations in the $C^{1+\mathrm{bv}}$-topology. We characterize the obstruction to minimality and describe the limit set. In particular, there are no invariant minimal Cantor sets, which can be seen as a Denjoy/Duminy type theorem for iterated systems on the circle.

Dédié à G. Duminy

1. Introduction

The rotation number is a classical tool in the study of dynamics of a single diffeomorphism of the circle. When the rotation number is rational, the map has periodic points and if it is irrational then either each orbit is dense on the circle or there is an invariant (minimal) Cantor set. Consider now a group of orientation-preserving homeomorphisms of the circle. The above classification can be extended to the group case where the orbit of a point is the group action on this point. Then, there can occur only one of the following three options [5]: existence of a finite orbit, every orbit is dense on the circle, or there exists a unique minimal Cantor set invariant by the group action. Minimal here means that all the points in the invariant set have dense orbits.

For iteration of single diffeomorphisms, the well-known theorem by Denjoy [3] implies that in the C^2-topology there cannot exist invariant minimal Cantor sets, while there are counter-examples in the C^1 class. Motivated by the study of co-dimension one foliations, Duminy extended the results of Denjoy for group actions [4]. Under the extra assumptions that the group generators are close to rotations in the $C^{1+\mathrm{bv}}$-topology (i.e. the C^1-class with bounded variation derivatives) and *finiteness of periodic points* for one of the generators, the invariant minimal Cantor set does not exist. See [13, 12] for details of the original proof.

In this paper, motivated by the dynamics of partially hyperbolic skew-products [6, 2, 11, 1], we study actions of finitely generated semigroups of diffeomorphisms on the circle, which can be viewed as iterated function systems. We extend the theorems of Denjoy and Duminy to the semigroup case and describe the limit set of possible orbits.

In order to state the main results, first a couple of definitions.

2010 *Mathematics Subject Classification.* Primary 58F15, 58F17; Secondary: 53C35.

Key words and phrases. Iterated Function Systems, invariant minimal Cantor sets, low-dimensional dynamics.

An *iterated function system* (IFS), generated by a finite family of diffeomorphisms $\Phi = \{\phi_1, \dots, \phi_k\}$ on the circle S^1, is the set $\mathrm{IFS}(\Phi)$ of all possible finite compositions of diffeomorphisms $\phi_i \in \Phi$. That is, the semigroup generated by the compositions of $\phi_1, \dots, \phi_k$.

The *orbit* of x for $\mathrm{IFS}(\Phi)$ is the action of IFS over the point x, i.e.,

$$\mathrm{Orb}^+_\Phi(x) \stackrel{\text{def}}{=} \{h(x) : \, h \in \mathrm{IFS}(\Phi)\} \subset S^1.$$

Denote by $\mathrm{Per}(\mathrm{IFS}(\Phi))$ the *set of periodic points* of $\mathrm{IFS}(\Phi)$, which is the set of $x \in M$ such that $h(x) = x$ for some $h \in \mathrm{IFS}(\Phi)$. We say the circle S^1 is minimal for $\mathrm{IFS}(\Phi)$ if the orbit of every point is dense.

Let f be a $C^{1+\mathrm{bv}}$-diffeomorphism on the circle. From the Denjoy Theorem the following statements are equivalent: S^1 is minimal for $\mathrm{IFS}(f)$ and f does not have periodic points. When the number of generators of the IFS increases the periodic points are no longer the obstruction to minimality. Now, this role is played by the ss-intervals which are compact intervals whose endpoints are consecutive attracting fixed points of different generators of $\mathrm{IFS}(\Phi)$. See Definition 2.1 for a formal definition.

Theorem A (Obstruction to minimality)**.** *There exists $\varepsilon > 0.38$ such that if f_0 and f_1 are diffeomorphisms of the circle with periodic points of period n_0 and n_1 respectively, ε-close to the rotations in the $C^{1+\mathrm{bv}}$-topology and with no periodic points in common, then the following conditions are equivalent:*

- *S^1 is minimal for $\mathrm{IFS}(f_0^{n_0}, f_1^{n_1})$,*
- *S^1 is minimal for $\mathrm{IFS}(f_0^{-n_0}, f_1^{-n_1})$,*
- *there are no ss-intervals for $\mathrm{IFS}(f_0^{n_0}, f_1^{n_1})$.*

When this occurs, both the hyperbolic attracting and the hyperbolic repelling periodic points of $\mathrm{IFS}(f_0^{n_0}, f_1^{n_1})$ are dense in S^1.

Under the same assumptions of Theorem A, the first theorem of Duminy [13, 12] can be restated as, S^1 is minimal for the group action and the hyperbolic periodic points are dense. See Theorem 5.4 in this article.

If we assume the generators have only hyperbolic periodic points, then Theorem A is robust in the following sense. There exists an open set $\mathcal{U} \subset \mathrm{Diff}^{1+\mathrm{bv}}(S^1) \times \mathrm{Diff}^{1+\mathrm{bv}}(S^1)$ with $(f_0, f_1) \in \mathcal{U}$ such that for every pair $(g_0, g_1) \in \mathcal{U}$ it holds that S^1 is minimal for $\mathrm{IFS}(g_0^{n_0}, g_1^{n_1})$ if and only if there are no ss-intervals for $\mathrm{IFS}(f_0^{n_0}, f_1^{n_1})$. Note that examples of robustly minimal IFS on the circle were given in [6, 7, 10]. Actually, the above robustness of Theorem A is valid for perturbations of (f_0, f_1) in the C^0-topology, i.e., in $\mathrm{Hom}(S^1) \times \mathrm{Hom}(S^1)$, but this will be taken up in a future work.

The existence of ss-intervals and the non-minimality of the circle leads us to attempt in comprehending the indecomposable pieces of global dynamics. Thus, we introduce the notion of a limit set for the orbits of an IFS.

The *ω-limit of $x \in S^1$ for* IFS(Φ) is the set

$$\omega_\Phi(x) \overset{\text{def}}{=} \{y : \exists\, (h_n)_n \subset \text{IFS}(\Phi) \text{ such that } \lim_{n\to\infty} h_n \circ \cdots \circ h_1(x) = y\},$$

while the *ω-limit of* IFS(Φ) is

$$\omega(\text{IFS}(\Phi)) \overset{\text{def}}{=} \text{cl}\big(\{y : \ \exists\, x \in M \text{ such that } \ y \in \omega_\Phi(x)\}\big),$$

where "cl" denotes the closure of a set. Similarly, the *backward* or *α-limit of* IFS(Φ) is defined as $\alpha(\text{IFS}(\Phi)) \overset{\text{def}}{=} \omega(\text{IFS}(\Phi^{-1}))$ where $\Phi^{-1} = \{\phi_1^{-1}, \dots, \phi_k^{-1}\}$. From the backward and forward limit, we define the *limit set of* IFS(Φ) as

$$L(\text{IFS}(\Phi)) \overset{\text{def}}{=} \omega(\text{IFS}(\Phi)) \cup \alpha(\text{IFS}(\Phi)).$$

The following result, in particular, shows the spectral decomposition of the limit set of IFS generated by Morse-Smale diffeomorphisms of the circle. That is, the limit set can be written as a finite union of disjoint (closed) maximal transitive sets. A (not necessarily IFS invariant) set $\Lambda \subset S^1$ is *transitive* for IFS(Φ) if

$$\Lambda \subset \overline{\text{Orb}^+_\Phi(x)} \quad \text{for some } x \in \Lambda.$$

The transitive set is *maximal* if there is no other transitive set Ω with $\Lambda \subset \Omega$.

Theorem B (Spectral decomposition)**.** *There exists $\varepsilon > 0.30$ such that, under the assumptions of Theorem A and if S^1 is not minimal for* IFS($f_0^{n_0}, f_1^{n_1}$)*, then*

$$L(\text{IFS}(f_0^{n_0}, f_1^{n_1})) = \bigcup K_i \quad \textit{and} \quad K_i \subset \overline{\text{Per}(\text{IFS}(f_0^{n_0}, f_1^{n_1}))}$$

where each K_i is a compact, maximal transitive set for IFS($f_0^{n_0}, f_1^{n_1}$)*. Moreover, each compact set K_i is either*

- *a single fixed point of $f_0^{n_0}$ of $f_1^{n_1}$ or*
- *a $**$-interval for* IFS($f_0^{n_0}, f_1^{n_1}$) *with $** \in \{ss, su, uu\}$.*[1]

In particular, if f_0 and f_1 are Morse-Smale diffeomorphisms then the above union is finite and the sets K_i are pairwise disjoints.

Observe that the spectral decomposition of Theorem B is given for the IFS generated by $f_0^{n_0}$ and $f_1^{n_1}$. It remains a question to describe the possible limit set of IFS(f_0, f_1), and if there is a spectral decomposition what are the pieces. An extension of the problem is to prove a spectral decomposition type result when the generators are far from rotations. Observe that in this case Cantor sets may appear in the decomposition.

With respect to dynamical decomposition of IFS from the ergodic perspective, let us mention the works of [9] and [15], where is shown the existence of finitely many stationary or SRB measures for step skew-products over the interval or the circle.

---

[1] For the precise definition see Definition 2.1.

A set Λ is (forward) invariant for IFS(Φ) if $\phi_i(\Lambda) \subset \Lambda$ for all $i = 1, \dots, k$. We say that a closed invariant set Λ is *minimal* for IFS(Φ) if every point of Λ has dense orbit in Λ or equivalently if

$$\Lambda = \phi_1(\Lambda) \cup \cdots \cup \phi_k(\Lambda) = \overline{\mathrm{Orb}^+_\Phi(x)} \quad \text{for all } x \in \Lambda.$$

Similarly to the case of group actions, a closed invariant minimal set Λ for IFS has to satisfy one of the following: Λ is a finite orbit, has non-empty interior, or is a Cantor set, see Theorem 5.2. Observe that apriori invariant minimal Cantor sets can exist in the pieces of the spectral decomposition of Theorem B. The next result shows that this cannot occur and can be thought of as a Denjoy-type result for the semigroup action.

Theorem C (Denjoy for IFS)**.** *There exists $\varepsilon > 0.30$ such that under the assumptions of Theorem A and if S^1 is not minimal, the only closed invariant minimal sets for* IFS$(f_0^{n_0}, f_1^{n_1})$ *are the ss-intervals. Moreover, there are no invariant minimal Cantor sets for* IFS(f_0, f_1).

In the case of iterations of a single diffeomorphism on the circle, the Denjoy examples [8] show the existence of invariant minimal Cantor sets in the C^1-topology. Thus one can expect that the above Theorem C fails in the C^1-topology. The recent work of Shinohara [14] points in this direction, showing that there is a closed invariant minimal set different from a ss-interval for IFS(f_0, f_1) with f_0 and f_1 close to the identity in the C^1-topology.

The paper is organized as follows. Firstly, in Section 2 we state and give a proof of a relevant result of Duminy. This is necessary in order to understand the next section, which is a generalization of Duminy's Theorem using a new notion of cycles for IFS and concludes with the proof of Theorem A. In Section 4 we show a Spectral Decomposition Theorem on the real line, and afterwards prove Theorem B. The trichotomy on the shape of invariant minimal sets is taken up in the next section and finally Theorem C and Duminy's Theorem under the assumptions of Theorem A are shown.

Notation: In what follows we will use the following notation:

$$\Phi = (f_0, f_1), \qquad \Phi^{-1} = (f_0^{-1}, f_1^{-1}), \qquad \Phi^n = (f_0^{n_0}, f_1^{n_1}), \qquad \Phi^{-n} = (f_0^{-n_0}, f_1^{-n_1}).$$

2. A RESULT OF DUMINY

In this section we will show an important result of Duminy which can be easily deduced from [12, 13]. In order to do a self-contained paper and to indicate the techniques used to derive the stronger result in Section 3, the complete proof is given. Before that, the formal definition of $**$-intervals is introduced.

Definition 2.1 ($**$-intervals)**.** *Consider f_0, f_1 orientation preserving homeomorphims on the real line. Let $[a, b]$ be a compact interval such that*

$$\mathrm{Fix}(f_i) \cap (a, b) = \emptyset \quad \text{and} \quad [a, b] \subset f_0([a, b]) \cup f_1([a, b]).$$

We say that $[a, b]$ is a $**$-interval for IFS(Φ) with $** \in \{ss, su\}$ when a and b satisfy the additional properties (see Figure 1):



Figure 1. Examples of $**$-intervals

- ss-interval*: a and b are, respectively, attracting fixed points of the restriction of f_0 and f_1 to $[a, b]$ and $f_0(b) \neq b$, $f_1(a) \neq a$.*
- su-interval*: a and b are either, an attractor-repeller or a repeller-attractor pair for the same map restricted to $[a, b]$, say an attractor-repeller pair for f_0. In this case, we ask that $f_1 > \mathrm{id}$ in $[a, b]$ and $f_1([a, b]) \cap (a, b) \neq \emptyset$.*

A uu-interval *for* $\mathrm{IFS}(\Phi)$ *is a ss-interval for* $\mathrm{IFS}(\Phi^{-1})$*. An unbounded interval* $[a, \infty]$ *is a*

- *s-interval for* $\mathrm{IFS}(\Phi)$*: if a is an attracted fixed point of the restriction of a map to $[a, \infty)$, say f_0, satisfying $f_0 < \mathrm{id}$ in (a, ∞) and $f_1 > \mathrm{id}$ in $[a, \infty)$.*

A u-interval *for* $\mathrm{IFS}(\Phi)$ *is a s-interval for* $\mathrm{IFS}(\Phi^{-1})$*. Analogously, the unbounded case $[-\infty, b]$ is defined in the same manner.*

Let I be an interval on the real line or on S^1. Consider an orientation preserving C^1-map f of I such that $Df(x) \neq 0$ for all $x \in I$. The non-negative number

$$\mathrm{Dist}(f, I) = \sup_{x,y \in I} \log \frac{Df(x)}{Df(y)}$$

is called *distortion constant* of f in I or *total variation* of $\log Df$ in I. We say that f belongs to the $C^{1+\mathrm{bv}}$ class on I if it has a bounded distortion constant in I. Observe that, by means of the mean-value theorem, C^2-maps with non-zero first derivative are $C^{1+\mathrm{bv}}$.

It is not difficult to see that every orientation preserving $C^{1+\mathrm{bv}}$-map f on the circle satisfies

$$e^{-V_f} \leq Df(x) \leq e^{V_f} \quad \text{and} \quad |f(x) - (x + \rho(x))| < 1 - e^{-V_f} \quad \text{for all } x \in S^1$$

where $\rho(f)$ is the rotation number of f and $V_f = \mathrm{Dist}(f, S^1)$ (see [12] for details). Therefore, if the distortion constant of f is small enough, then f is close to a rotation. We will say that f is *ε-close to rotation in the $C^{1+\mathrm{bv}}$-topology* if $V_f \leq \varepsilon$.

Let f_0 and f_1 be orientation preserving $C^{1+\mathrm{bv}}$-diffeomorphisms on the real line. In what follows, we will study IFS(Φ) with f_0 and f_1 restricted to a $**$-interval $K^{**}=[a,\,b]$ for $**\in\{ss,s,su\}$. In the case of uu-intervals and u-intervals for IFS(Φ) the same results follow for IFS(Φ^{-1}). Observe that

$$K^{**}=f_0(K^{**})\cup f_1(K^{**}) \quad \text{for } **\in\{ss,s\}$$

and then, it is an invariant set by the action of IFS(Φ). However the above equality does not follow for a su-interval.

We will be interested in stating conditions under which every point in a $**$-interval has a dense orbit for IFS(Φ). In this case of a su-interval and a s-interval notice that one of the endpoints of K^{su} and K^{s} can never have a dense orbit. To unify the notation, we write

$$K^{**}\subset\overline{\mathrm{Orb}^{+}_{\Phi}(x)} \quad \text{for all } x\in K^{**}$$

and in this case say the $**$-interval is *minimal* for IFS(Φ).

Theorem 2.2 (Duminy's Lemma)**.** *Assume* $f_0<\mathrm{id}$ *in* (a,b) *and suppose that there exists* $\varepsilon>0$ *such that*

$$|Df_0(x)-1|<\varepsilon \quad \textit{for all } x\in(a,\,f_0^{-1}f_1(a)) \quad \textit{and} \quad (1-\varepsilon)\varepsilon^{-1}e^{-V}>1$$

where $V=(V_0+V_1)|a-f_0^{-1}f_1(a)|$ *being* V_0 *and* V_1 *constants satisfying*

$$\mathrm{Dist}(f_0,I)\le V_0 \quad \textit{and} \quad \mathrm{Dist}(f_1,J)\le V_1$$

for all fundamental domain I *of* f_0 *and* J *of* f_1 *in* $(a,\,f_0^{-1}f_1(a)]$. *Then*

$$K^{**}\subset\overline{\mathrm{Per}(\mathrm{IFS}(\Phi))} \quad \textit{and} \quad K^{**}\subset\overline{\mathrm{Orb}^{+}_{\Phi}(x)} \ \ \textit{for all } x\in K^{**}.$$

Notice that from definition of $**$-intervals for IFS(Φ) with $**\in\{ss,su,s\}$, the *overlap condition* is verified, that is, $f_0(K^{**})\cap f_1(K^{**})\neq\emptyset$. This condition implies that

$$A=(f_1(a),\,f_0^{-1}f_1(a)]\subset[a,\,b].$$

Next, we will define a first return map $\mathcal{R}$ over the fundamental domain A.

For each $x\in A$ let $m(x)\ge 1$ be the smallest positive number such that $f_1^{-m(x)}(x)\notin A$ and let $n(x)$ be the first time for which $f_1^{-m(x)}(x)$ returns to A by iterations of f_0^{-1}. Then we can define the first return map $\mathcal{R}$ in the following way

$$\mathcal{R}\colon A\to A, \qquad \mathcal{R}(x)=f_0^{-n(x)}f_1^{-m(x)}(x).$$

Note that this map can be written as $\mathcal{R}(x)=F^{n(x)+m(x)}(x)$ for $x\in A$, where $F:[a,\,b]\to[a,\,b]$ is defined by $F=f_0^{-1}$ in $[a,\,f_1(a)]$ and $F=f_1^{-1}$ in $(f_1(a),\,b]$. Therefore, for every $x\in[a,\,b]$ there is a smallest non-negative number $k(x)\ge 0$ such that $F^{k(x)}(x)\in A$ and $\mathcal{R}$ can be extended to the whole interval by taking

$$\mathcal{R}\colon(a,b)\to A, \qquad \mathcal{R}(x)=F^{n+m+k}(x),$$

where $k=k(x)$, $m=m(F^k(x))$ and $n=n(F^k(x))$.

A point $d \in A$ is said to be a *discontinuity of* $\mathcal{R}$ if $\mathcal{R}(d) = f_0^{-1}f_1(a)$ or equivalently, if $d = f_1^{m(d)}f_0^{n(d)-1}f_1(a)$. These points define a partition on A. In other to describe this partition we have to consider two cases: $f_1^2(a) \notin A$ and $f_1^2(a) \in A$. In the first case $m(x) = 1$ for all $x \in A$ and we write $I_1 = A$. In the second case, consider $\ell \in \mathbb{N}$ such that $f_1^\ell f_1(a) \in A$, but $f_1^{\ell+1}f_1(a) \notin A$. Then $f_1^m f_1(a)$ for $m = 1, 2, \ldots, \ell$ defines a partition on A given by

$$\begin{aligned} I_\ell &= (f_1^\ell f_1(a),\, f_0^{-1}f_1(a)] \quad \text{and} \\ I_m &= (f_1^m f_1(a),\, f_1^m f_1(a)] \quad \text{for } 1 \le m < \ell. \end{aligned}$$

Hence, $m(x) = m$ for each $x \in I_m$ and $f_1^{-m(x)}(I_m) \subset (a,\, f_1(a)]$. Observe that the sequence of points $f_0^n f_1(a)$, $n \ge 0$, creates a partition on the interval $(a,\, f_1(a)]$. This partition induces a partition in right-closed intervals I_{mn} in each I_m. Namely, $I_{mn} \subset f_1^m f_0^n(A)$ for $1 \le m \le \ell$ and $n \ge 1$.

Proposition 2.3. *For every $J = [x, y] \subset I_{mn}$ it holds that*

$$\frac{|\mathcal{R}(J)|}{|J|} \ge (1-\varepsilon)\varepsilon^{-1}e^{-V}\,\frac{D(\mathcal{R}(y))}{D(y)}$$

where $D : A \to (0, \infty)$ is given by $D(y) = |f_0(y) - y|$.

Proof. Notice that $\mathcal{R}(J) = f_0^{-n}f_1^{-m}(J)$ and hence

$$\frac{|\mathcal{R}(J)|}{|J|} = \frac{|f_0^{-n}f_1^{-m}(J)|}{|f_1^{-m}(J)|} \cdot \frac{|f_1^{-m}(J)|}{|J|}. \tag{1}$$

In order to estimate the first factor, observe that $f_1^{-m}(J) \subset [f_0 f_1^{-m}(y),\, f_1^{-m}(y)]$ and so

$$\begin{aligned} \mathrm{Dist}(f_0^{-n},&[f_0f_1^{-m}(y),\, f_1^{-m}(y)]) = \mathrm{Dist}(f_0^n, [f_1^{-m}(y),\, f_0^{-1}f_1^{-m}(y)]) \\ &\le \sum_{i=0}^{n-1} \mathrm{Dist}(f_0, f_0^i([f_1^{-m}(y), f_0^{-1}f_1^{-m}(y)])) \le V_0|a - f_0^{-1}f_1(a)|. \end{aligned}$$

Thus,

$$\frac{|f_0^{-n}f_1^{-m}(J)|}{|f_0^{-n}f_0f_1^{-m}(y) - f_0^{-n}f_1^{-m}(y)|} \ge e^{-V_0|a-f_0^{-1}f_1(a)|}\,\frac{|f_1^{-m}(J)|}{|f_0f_1^{-m}(y) - f_1^{-m}(y)|}. \tag{2}$$

Let us denote by $t = f_1^{-m}(y)$. Using the mean value theorem, there is some $\xi \in (a, t)$ such that $f_0(t) - a = f_0(t) - f_0(a) = Df_0(\xi)(t - a)$. Hence

$$f_0(t) - t = \frac{Df_0(\xi) - 1}{Df_0(\xi)}\,(f_0(t) - a) \le \frac{\varepsilon}{1-\varepsilon}\,(f_0(t) - a).$$

Substituting in (2) one has that

$$\frac{|f_0^{-n}f_1^{-m}(J)|}{|f_1^{-m}(J)|} \ge (1-\varepsilon)\varepsilon^{-1}e^{-V_0|a-f_0^{-1}f_1(a)|}\,\frac{|f_0(\mathcal{R}(y)) - \mathcal{R}(y)|}{|f_0f_1^{-m}(y) - a|}. \tag{3}$$

Now we will estimate the second factor in (1). Observe that $f_1^{-m}(J)$ and $[a,\, f_1^{-m}(y)]$ are contained in the fundamental domain $[a,\, f_1(a)]$ of f_1. Thus,

$$\mathrm{Dist}(f_1^m, [a,\, f_1(a)]) \le \sum_{i=0}^{m-1} \mathrm{Dist}(f_1, f_1^i([a,\, f_1(a)])) \le V_1|a - f_0^{-1}f_1(a)|$$

and so

$$\frac{|J|}{|f_1^m(a)-y|}=\frac{|f_1^m f_1^{-m}(J)|}{|f_1^m([a,\,f_1^{-m}(y)])|}\leq e^{V_1|a-f_0^{-1}f_1(a)|}\,\frac{|f_1^{-m}(J)|}{|f_1^{-m}(y)-a|}.$$

As $a<f_0f_1^{-m}(y)<f_1^{-m}(y)$ and since $y\in I_{mn}$, by the construction of the partition I_{mn} of I_m, one has that $f_0(y)\leq f_1(a)\leq f_1^m(a)<y$. Then the above inequality implies

$$\frac{|f_1^{-m}(J)|}{|J|}\geq e^{-V_1|a-f_0^{-1}f_1(a)|}\,\frac{|f_0f_1^{-m}(y)-a|}{|f_0(y)-y|} \tag{4}$$

Putting together (3) and (4) in Equation (1) one deduces the inequality desired and concludes the proof of the lemma. □

The following lemma shows that some iterate of the first return map $\mathcal{R}$ is expanding.

Lemma 2.4. *There exists $N\in\mathbb{N}$ such that $D\mathcal{R}^N(x)>1$ for all $x\in A$.*

Proof. Let us denote by

$$C(f_0)=\frac{\inf\{D(x):x\in A\}}{\sup\{D(x):x\in A\}}>0.$$

Fix $\kappa>1$ and let $N=N(f_0)$ be a natural number such that

$$((1-\varepsilon)\varepsilon^{-1}e^{-V})^N\,C(f_0)>\kappa.$$

Suppose $x\in I_{mn}$ and consider $\delta>0$ small enough such that $J=[x-\delta,x]\subset I_{mn}$. Hence, by Lemma 2.3 it follows that

$$\begin{aligned}\frac{|\mathcal{R}^N(J)|}{|J|}&=\frac{|\mathcal{R}^N(J)|}{|\mathcal{R}^{N-1}(J)|}\cdots\frac{|\mathcal{R}(J)|}{|J|}\\&\geq((1-\varepsilon)\varepsilon^{-1}e^{-V})^N\,\frac{D(\mathcal{R}^N(x))}{D(x)}\\&\geq((1-\varepsilon)\varepsilon^{-1}e^{-V})^N\,C(f_0)>\kappa.\end{aligned}$$

Since this inequality holds for every $\delta>0$ small enough, this allows to conclude that $D\mathcal{R}^N(x)\geq\kappa>1$. □

The first simplification to prove the minimality and the density of periodic points in a $**$-interval is to note that it suffices to show that the orbit of the global attractor a for $f_0|_{K^{**}}$, with respect to for IFS(Φ), is dense in the interval $K^{**}=[a,b]$.

Lemma 2.5. *If $K^{**}\subset\overline{\mathrm{Orb}^+_\Phi(a)}$ then*

$$K^{**}\subset\overline{\mathrm{Per}(\mathrm{IFS}(\Phi))}\quad\textit{and}\quad K^{**}\subset\overline{\mathrm{Orb}^+_\Phi(x)}\quad\textit{for all } x\in K^{**}.$$

Proof. Consider $x\in K^{**}$ and let V be any open set in K^{**}. From the density of the orbit of a for IFS(Φ) in K^{**}, there is $h\in\mathrm{IFS}(\Phi)$ such that $h(a)\in V$. Since h is a continuous map and a is a global attractor of f_0 in K^{**}, there exists $\ell\in\mathbb{N}$ such that $f_0^\ell(x)$ is close enough to 0, so that $h\circ f_0^\ell(x)\in V$. Therefore, the orbit of x for IFS(Φ) is dense in K^{**}.

Now, given $x\in K^{**}$ we will show that $x\in\overline{\mathrm{Per}(\mathrm{IFS}(\Phi))}$. Let I be any open interval such that $x\in I$. From the density of the orbit of a, there is $h\in\mathrm{IFS}(\Phi)$ such that $h(a)\in I$. Since

h is a continuous map there is $\delta > 0$ such that $h((a-\delta,\, a+\delta)) \subset I$. Using that a is a global attractor of f_0, there is $\ell \geq 0$ such that $f_0^\ell(I) \subset (a-\delta,\, a+\delta)$. Then $h \circ f_0^\ell(I) \subset I$. By the Banach fixed point theorem, $h \circ f_0^\ell$ has a (unique) attracting fixed point in I and thus $I \cap \mathrm{Per}(\mathrm{IFS}(\Phi)) \neq \emptyset$. This implies that x belongs to the closure of the periodic points of $\mathrm{IFS}(\Phi)$ and we conclude the proof of the lemma. ☐

Now, we are ready to proof Theorem 2.2

Proof of Theorem 2.2. Recall that the return map $\mathcal{R} : A \to A$ can be extended to the interval (a, b). In particular, this implies that for any open interval $I \subset K^{**}$, there exists $h \in \mathrm{IFS}(\Phi)$ such that $h^{-1}(I) \cap A \neq \emptyset$. From Lemma 2.4, the return map $\mathcal{R}^N$ is an expanding map in A. Thus, there is $\ell \in \mathbb{N}$ such that $\mathcal{R}^{\ell N}(h^{-1}(I) \cap A)$ contains some discontinuity of $\mathcal{R}^N$.

Recall that the discontinuities $d \in A$ are points in the orbit of a for $\mathrm{IFS}(\Phi)$, i.e., applying the appropriate inverse branch, $d = \mathcal{R}^{-N}(f_0^{-1} f_1(a)) = L \circ f_1(a)$ for some $L \in \mathrm{IFS}(\Phi)$. Then, one has $h \circ L \circ f_1(a) \in I$. Therefore, the orbit of a is dense in K^{**} with respect to $\mathrm{IFS}(\Phi)$. Finally, from Lemma 2.5 we conclude the proof of the theorem. ☐

3. Cycles for IFS and proof of Theorem A

Now we will give a proof of Theorem A. We will first show the result when the rotation number of f_0 and f_1 is zero, that is, both of the diffeomorphisms have fixed points. From now on assume that there are no ss-intervals for $\mathrm{IFS}(\Phi)$, and so we have to show minimality of S^1. The proof will require the following notion of a cycle for an IFS of two diffeomorphisms on the circle. Similarly to Duminy's result (Theorem 2.2), using the cycle we will then construct a return map and afterwards estimate the derivative. Here the expanding return map is of a global character, while Duminy's lemma can be thought of as a local version.

3.1. Cycles.

Since f_0 and f_1 do not have fixed points in common, we can (without ambiguity) denote by $\{s_i\}$ the set of fixed points of f_0 and f_1 with a non-empty basin of attraction. This set includes the fixed points which may attract only from one-side. Let $B(s_i)$ represent the basin of attraction of s_i with respect the relevant map f_0 or f_1 which does not include the point s_i (it is an open set).

Definition 3.1 (Cycle for IFS). *Define a partial order on the attracting points by $s_i \prec s_j$ if and only if s_i belongs to $B(s_j)$. A sequence of attractors is said to be a* cycle of length n for $\mathrm{IFS}(\Phi)$ *if*

$$s_{i_1} \prec s_{i_2} \prec \cdots \prec s_{i_n} \prec s_{i_{n+1}} \quad \textit{and} \quad s_{i_1} = s_{i_{n+1}}.$$

Notice that:

- since the attracting points s_i of the cycle alternate with respect to the maps f_0 and f_1, the length of a cycle is always even.
- a ss-interval is a cycle of length 2.

The following result shows the existence of a cycle under the assumption of no periodic points in common.

Proposition 3.2. *Let f_0, f_1 be circle diffeomorphisms with zero rotation number and no fixed points in common. Then there exists at least one cycle for* IFS(Φ)*.*

Proof. Since f_0 and f_1 have no fixed points in common, given $x \in S^1$, then $x \in B(s_i)$ for some s_i. By compactness we can take a finite sub-covering and re-ordering the indices, we may suppose that $S^1 = B(s_1) \cup \cdots \cup B(s_n)$.

For any s_i, there exists $s_{i_2} \in \{s_1, \ldots, s_n\}$ with $s_i \prec s_{i_2}$. Proceeding inductively we obtain the sequence $(s_i = s_{i_1})$, $s_{i_1} \prec s_{i_2} \prec \cdots \prec s_{i_n} \prec s_{i_{n+1}}$. Then necessarily $s_{i_{n+1}} = s_{i_j}$ with $j \in \{1, \ldots, n\}$, and so we have the cycle,

$$s_{i_j} \prec s_{i_2} \prec \cdots \prec s_{i_n} \prec s_{i_j}. \qquad \square$$

By the above proposition we may assume that there exists a cycle of length n for IFS(Φ) and re-numbering the indices, the cycle can be written as

$$s_n \prec s_{n-1} \prec \cdots \prec s_2 \prec s_1 \prec s_0 \quad \text{with} \quad s_0 = s_n,$$

where s_0 is a (semi)attracting fixed point of f_0. Note that s_k is a (semi)attracting fixed point for $f_{k \bmod 2}$: if k is an even number then s_k is a fixed point of f_0 and if k is an odd number then of f_1.

Notation: For the rest of the section f_k will stand for $f_{k \bmod 2}$.

We consider S^1 parameterized by $[s_0,\, s_0 + 1] \bmod 1$, and so $s_0 < s_k < s_0 + 1 = s_n$ for $k = 1, \ldots, n-1$ with respect to the real order on the interval $[s_0,\, s_0 + 1]$. We denote by s_k^- and s_k^+ (when they exist) the fixed points of the lift of f_k on the real line closest to s_k from the left and right, respectively. If f_k has more than one fixed point and since s_k is a (semi)-attractor, necessarily one of the points, s_k^- or s_k^+, exists and is a (semi)-repeller. With respect to the order on the interval $[s_0,\, s_0 + 1]$, and as $s_1 \prec s_0$, we may suppose that $s_1 \in (s_0,\, s_0^+) \subset B(s_0)$. The next lemma on the order of the points is crucial for the creation of the return map. The combinatorics for the other case when $s_1 \in (s_n^-,\, s_n) \subset B(s_n)$ are different but the construction of the return map is easily adapted.

Lemma 3.3. *If there are no ss-intervals for* IFS(Φ) *then on the interval* $[s_0,\, s_0 + 1]$ *the cycle has the order*

$$s_0 < s_1 < \cdots < s_{n-1} < s_0 + 1 = s_n, \quad \textit{where } s_{k+1} \prec s_k \ \textit{ for } \ k = 0, \ldots, n-1.$$

Moreover,

$$s_k < f_{k+1}^{-1}(s_k) < s_{k+1} < s_k^+ \quad \textit{for } k = 0, \ldots, n-1.$$

Proof. We will show that $s_{k+1} \in (s_k,\, s_k^+) \subset B(s_k)$ for every $k = 0, \ldots, n-1$ and this proves the first part of the lemma. The second part will follow as the fixed point of f_{k+1} closest to s_k from the right has to be a repeller from the left (otherwise a ss-interval is created) and so $s_k < f_{k+1}^{-1}(s_k) < s_{k+1} < s_k^+$.

Firstly, we will assume that f_0 or f_1 has a unique fixed point. In that case, the cycle has length 2 and since there are no ss-intervals then $s_2 = s_0+1$. Thus $s_0 < s_1 < s_0^+ \leq s_0+1 = s_2$, where $s_2 \prec s_1 \prec s_0$. In particular, $s_1 \in (s_0,\, s_0^+) \subset B(s_0)$ and $s_2 \in (s_1,\, s_1^+) \subset B(s_1)$.

Now, we will assume that f_0 and f_1 have more than one fixed point. By the initial hypothesis $s_1 \in (s_0,\, s_0^+) \subset B(s_0)$, and let us show that $s_2 \in (s_1,\, s_1^+) \subset B(s_1)$. Since $s_2 \prec s_1$, $s_2 \in B(s_1) \subset (s_1^-,\, s_1) \cup (s_1,\, s_1^+)$ and so or $s_2 \in (s_1^-,\, s_1)$ or $s_2 \in (s_1,\, s_1^+)$. It is enough to prove that the first case cannot occur.

Suppose that $s_2 \in (s_1^-,\, s_1)$. If $s_0 < s_1^-$ then $s_0 < s_2 < s_1 < s_0^+$, which is a contradiction since s_0^+ is the closest fixed point of f_0 to s_0 from the right. In the other case, $s_0 > s_1^-$, a similar argument concludes that $s_0 = s_2$. Since there are no ss-intervals and $s_2 \prec s_1$ then $B(s_1) = (s_1,\, s_1^+)$ and $s_0 + 1 = s_2 + 1 < s_1^+$. Thus $s_1^- < s_0 < s_0 + 1 < s_1^+$, which implies f_1 has a unique fixed point, a contradiction. Therefore s_2 does not belong to $(s_1^-,\, s_1)$ and so it is contained in $B(s_1)$.

Suppose by the inductive hypothesis that $s_i \in (s_{i-1},\, s_{i-1}^+) \subset B(s_{i-1})$ for all $i \leq k$. To prove $s_{k+1} \in (s_k,\, s_k^+) \subset B(s_k)$ the idea is basically the same as above. If $s_{k+1} \in (s_k^-,\, s_k)$, then $s_k^- \in (s_{k-1},\, s_k)$ or $s_k^- < s_{k-1}$. The first case cannot occur because s_{k+1} and s_{k-1} are fixed points of the same map $f_{k+1} = f_{k-1}$ and $s_{k+1} \in (s_k^-,\, s_k)$. Thus $s_k^- < s_{k-1}$ and so $s_{k+1} \leq s_{k-1}$. This fact together with $s_{k+1} \prec s_k$ imply that $(s_k^-,\, s_k) \subset B(s_k)$ and hence $[s_{k-1},\, s_k]$ is a ss-interval, and thus we get a contradiction. Therefore $s_k < s_{k+1} < s_k^+$ and so $(s_k,\, s_k^+) \subset B(s_k)$, which completes the induction and concludes the proof of the lemma. □

3.2. **Creating a return map.** Let us now use the cycle from the previous section

$$s_n \prec s_{n-1} \prec \cdots \prec s_2 \prec s_1 \prec s_0$$

to create a return map. Remember, f_k stands for $f_{k \bmod 2}$.

Lemma 3.4. *There exist families of right-closed pairwise disjoint intervals* $I_{i_1\dots i_n} \subset S^1$ *and maps* $h_{i_i\dots i_n} \in \mathrm{IFS}(\Phi)$ *with* $i_j \geq 0$ *for* $j = 1, \dots, n$ *such that*

- *i)* $A = (s_0,\, f_1^{-1}(s_0)] = \bigcup I_{i_1\dots i_n}$,
- *ii)* $h^{-1}_{i_1\dots i_n}(I_{i_1\dots i_n}) \subset A$ *if* $i_n = 0$ *and* $h^{-1}_{i_1\dots i_n}(I_{i_1\dots i_n}) = A$ *if* $i_n > 0$,
- *iii) if* $c \in A \setminus \{f_1^{-1}(s_0)\}$ *is an endpoint of* $I_{i_1\dots i_n}$ *then there exist a map* $h \in \mathrm{IFS}(\Phi)$ *and a point* $s \in \mathrm{Orb}^+_\Phi(s_1) \cup \cdots \cup \mathrm{Orb}^+_\Phi(s_n)$ *such that* $h(s) = c$. *That is, the point* c *is in the orbit of the cycle for* $\mathrm{IFS}(\Phi)$.

Proof. We can define a non-empty fundamental domain of f_{k+1},

$$A_k = (s_k,\, f^{-1}_{k+1}(s_k)] \quad \text{for } k = 0, \dots, n.$$

Since on the circle $s_0 = s_n$, we identify on S^1 the intervals in the real line $A_0 = (s_0, f_1^{-1}(s_0)]$ and $A_n = (s_n, f_1^{-1}(s_n)]$, denoting this interval by A. In order to create the expanding return map we will divide this fundamental domain inductively.

By Lemma 3.3, $s_0 < f_1^{-1}(s_0) < s_1$ and since $s_1 \prec s_0$, there exists $j \in \mathbb{N}$ such that $s_0 < f_0^j(s_1) < f_1^{-1}(s_0) \leq f_0^{j-1}(s_1) < s_1$. Then

$$A = (s_0,\, f_1^{-1}(s_0)] = \bigcup_{i_1=0}^{\infty} I_{i_1},$$

with

$$I_0 = (f_0^j(s_1),\, f_1^{-1}(s_0)] \quad \text{and} \quad I_{i_1} = (f_0^{j+i_1}(s_1),\, f_0^{j+i_1-1}(s_1)] \ \text{ if } i_1 > 0.$$

Let $h_{i_1} = f_0^{j+i_1}$, we have that $h_{i_1}^{-1}(I_{i_1}) = (s_1,\, c_{i_1}]$ where

$$c_0 = f_0^{-j} f_1^{-1}(s_0) \in (s_1,\, f_0^{-1}(s_1)] \quad \text{and} \quad c_{i_1} = f_0^{-1}(s_1) \ \text{ if } i_1 > 0.$$

Therefore, $h_0^{-1}(I_0) \subset A_1$ and $h_{i_1}^{-1}(I_1) = A_1$ if $i_1 > 0$.

Let $c \in A \setminus \{f_1^{-1}(s_0)\}$ be an endpoint of I_{i_1}. Then, either, $c = f_0^j(s_1)$ if $i_1 = 0$ or $c \in \{f_0^{j+i_1}(s_1), f_0^{j+i_1-1}(s_1)\}$ if $i_1 > 0$. In any case, c belongs to the orbit of the cycle. This completes the first step of the induction and now we proceed with the inductive hypothesis.

Suppose that we have families of right-closed pairwise disjoint intervals $I_{i_1\ldots i_k} \subset S^1$ and maps $h_{i_1\ldots i_k} \in \mathrm{IFS}(\Phi)$ with $i_j \geq 0$ for $j = 1, \ldots, k$ such that

(i) $A = \bigcup I_{i_1\ldots i_k}$.

(ii) $h_{i_1\ldots i_k}^{-1}(I_{i_1\ldots i_k}) = (s_k,\, c_{i_1\ldots i_k}]$ where

$$c_{i_1\ldots i_{k-1}0} \in (s_k,\, f_{k+1}^{-1}(s_k)] \quad \text{and} \quad c_{i_1\ldots i_k} = f_{k+1}^{-1}(s_k) \ \text{ if } i_k > 0.$$

In particular, $h_{i_1\ldots i_{k-1}0}^{-1}(I_{i_1\ldots i_{k-1}0}) \subset A_k$ and $h_{i_1\ldots i_k}^{-1}(I_{i_1\ldots i_k}) = A_k$ if $i_k > 0$.

(iii) If $c \in A \setminus \{f_1^{-1}(s_0)\}$ is an endpoint of $I_{i_1\ldots i_k}$ then there exist a map $h \in \mathrm{IFS}(\Phi)$ and a point $s \in \mathrm{Orb}_\Phi^+(s_0) \cup \cdots \cup \mathrm{Orb}_\Phi^+(s_n)$ such that $h(s) = c$.

By Lemma 3.3 $s_k < f_{k+1}^{-1}(s_k) < s_{k+1} < s_k^+$. Hence, from the inductive hypothesis we get

$$s_k < c_{i_1\ldots i_k} \leq f_{k+1}^{-1}(s_k) < s_{k+1} < s_k^+.$$

Now, since $s_{k+1} \prec s_k$, for each multi-index $i_1 \ldots i_k$ there exists $j_{i_1\ldots i_k} \in \mathbb{N}$ such that

$$s_k < f_k^{j_{i_1\ldots i_k}}(s_{k+1}) < c_{i_1\ldots i_k} \leq f_k^{j_{i_1\ldots i_k}-1}(s_{k+1}) < s_{k+1}.$$

Then

$$h_{i_1\ldots i_k}^{-1}(I_{i_1\ldots i_k}) = (s_k,\, c_{i_1\ldots i_k}] = \bigcup_{\ell=0}^{\infty} J_{i_1\ldots i_k\ell}$$

with

$$J_{i_1\ldots i_k 0} = (f_k^{j_{i_1\ldots i_k}}(s_{k+1}),\, c_{i_1\ldots i_k}] \ \text{ and}$$
$$J_{i_1\ldots i_k\ell} = (f_k^{j_{i_1\ldots i_k}+\ell}(s_{k+1}),\, f_k^{j_{i_1\ldots i_k}+(\ell-1)}(s_{k+1})] \ \text{ if } \ell > 0.$$

Notice that the intervals $J_{i_1\ldots i_k\ell}$, $\ell \geq 0$ are pairwise disjoint. Define $I_{i_1\ldots i_k\ell} \subset I_{i_1\ldots i_k}$ by

$$I_{i_1\ldots i_k\ell} = h_{i_1\ldots i_k}(J_{i_1\ldots i_k\ell}).$$

By definition, for a fixed multi-index $i_1 \ldots i_k$, the intervals $I_{i_i\ldots i_k\ell}$, $\ell \geq 0$ are also pairwise disjoint. Hence, by means of the induction hypothesis on the intervals $I_{i_1\ldots i_k}$ and since

$I_{i_1\dots i_k\ell} \subset I_{i_1\dots i_k}$ it follows that $\{I_{i_1\dots i_{k+1}} : \ i_j \geq 0 \text{ for } j=1,\dots,k+1\}$ is a family of right-closed pairwise disjoint intervals. Note that each right-closed interval $I_{i_1\dots i_k}$ is union of the intervals $I_{i_1\dots i_k\ell}$, $\ell \geq 0$. Then, by the induction hypothesis it also follows that $A = \cup I_{i_1\dots i_{k+1}}$.

Set $h_{i_1\dots i_k i_{k+1}} = h_{i_1\dots i_k} \circ f_k^{j_{i_1\dots i_k}+i_{k+1}}$. By construction

$$h^{-1}_{i_1\dots i_{k+1}}(I_{i_1,\dots i_{k+1}}) = f_k^{-j_{i_1\dots i_k}-i_{k+1}}(J_{i_1\dots i_k i_{k+1}}) = (s_{k+1},\, c_{i_1\dots i_{k+1}}]$$

where $c_{i_1\dots i_k i_{k+1}} = f_k^{-1}(s_{k+1})$ if $i_{k+1} > 0$ and

$$c_{i_1\dots i_k 0} = h^{-1}_{i_1\dots i_k 0} \circ h_{i_1\dots i_k}(c_{i_1\dots i_k}) = f_k^{-j_{i_1\dots i_k}}(c_{i_1\dots i_k}) \in (s_{k+1},\, f_k^{-1}(s_{k+1})].$$

Therefore, $h^{-1}_{i_1\dots i_k 0}(I_{i_1\dots i_k 0}) \subset A_{k+1}$ and $h^{-1}_{i_1\dots i_{k+1}}(I_{i_1\dots i_{k+1}}) = A_{k+1}$ if $i_{k+1} > 0$.

In order to prove the third item, we fix an interval $I_{i_1\dots i_k\ell}$. For every $\ell \geq 0$ the left endpoint of this interval is $h_{i_1\dots i_k} \circ f_k^{j_{i_1\dots i_k}+\ell}(s_{k+1})$ and so it belongs to the orbit of the cycle. The right endpoint is

$$h_{i_1\dots i_k}(c_{i_1\dots i_k}) \ \text{ if } \ \ell = 0 \quad \text{and} \quad h_{i_1\dots i_k} \circ f_k^{j_{i_1\dots i_k}+\ell-1}(s_{k+1}) \ \text{ if } \ \ell > 0.$$

Note that $h_{i_1\dots i_k}(c_{i_1\dots i_k})$ is a endpoint of $I_{i_1\dots i_k}$. Therefore, by the inductive hypothesis, either $h_{i_1\dots i_k}(c_{i_1\dots i_k}) = f_1^{-1}(s_0)$ or $h_{i_1\dots i_k}(c_{i_1\dots i_k}) = h(s)$ for some $h \in \mathrm{IFS}(\Phi)$ and $s \in \cup_{i=0}^n \mathrm{Orb}^+_\Phi(s_i)$.

Going through the n steps of the cycle we conclude the lemma. □

From the above lemma we define the return map over $A = (s_0,\, f_1^{-1}(s_0)]$ as

$$\mathcal{R}\colon A \to A, \qquad \mathcal{R}|_{I_{i_1\dots i_n}} = h^{-1}_{i_1\dots i_n}.$$

The endpoints of the intervals $I_{i_1\dots i_n}$ are called *discontinuities* of $\mathcal{R}$ and if s is a discontinuity then $s \in \mathrm{Orb}^+_\Phi(s_0) \cup \dots \cup \mathrm{Orb}^+_\Phi(s_n)$. We summarize some of the points from the inductive process in the construction of the return map in Lemma 3.4. This notation will be required for the next section.

Addendum 3.5. *For each $k = 1,\dots,n$, there exists a family*

$$\{(I_{i_1\dots i_k}, h_{i_1\dots i_k}, m_{i_1\dots i_k}) : i_j \geq 0 \ j = 1,\dots k\}$$

with $I_{i_1\dots i_k}$ pairwise disjoint right-closed intervals of S^1, $h_{i_1\dots i_k} \in \mathrm{IFS}(\Phi)$ and $m_{i_1\dots i_k}$ natural numbers such that $A = (s_0,\, f_1^{-1}(s_0)] = \cup I_{i_1\dots i_k}$, and for $k = 1,\dots,n-1$

- *i) $I_{i_1\dots i_k i_{k+1}} \subset I_{i_1\dots i_k}$ and I_{i_1}, $h_{i_1\dots i_k}(I_{i_1\dots i_{k+1}})$ are respectively contained in a fundamental domain of f_0 and f_k;*
- *ii) $m_{i_1} = j + i_1$, $m_{i_1\dots i_k} = j_{i_1\dots i_k} + i_{k+1}$, and $h^{-1}_{i_1} = f_0^{-m_{i_1}}$,*

$$h^{-1}_{i_1\dots i_k i_{k+1}} = f_k^{-m_{i_1\dots i_{k+1}}} \circ h^{-1}_{i_1\dots i_k} = f_k^{-m_{i_1\dots i_{k+1}}} f_{k-1}^{-m_{i_1\dots i_k}} \cdots f_1^{-m_{i_1 i_2}} f_0^{-m_{i_1}};$$

- *iii) $h^{-1}_{i_1\dots i_{k-1}0}(I_{i_1\dots i_{k-1}0}) = (s_k,\, c_{i_1\dots i_k}]$ and $h^{-1}_{i_1\dots i_k}(I_{i_1\dots i_k}) = A_k$ if $i_k > 0$ with*

$$s_k < c_{i_1\dots i_k} \leq f^{-1}_{k+1}(s_k) \quad \textit{and} \quad A_k = (s_k,\, f^{-1}_{k+1}(s_k)];$$

- *iv) $m_{i_1\dots i_k+1} = m_{i_1\dots i_k} + 1$ and $m_{i_1\dots i_{k-1}0} \geq 1$ satisfies that*

$$f_k^{m_{i_1\dots i_{k-1}0}}(s_{k+1}) < c_{i_1\dots c_{i_k}} \leq f_k^{m_{i_1\dots i_{k-1}0}-1}(s_{k+1}).$$

3.3. **Estimation of the derivative for the return map.** We will prove the following proposition which can be compared to Proposition 2.3.

Proposition 3.6. *For every* $J = [x, y] \subset I_{i_1 \dots i_n}$ *it holds that*

$$\frac{|\mathcal{R}(J)|}{|J|} \geq \left(\frac{e^{-(V_0+V_1)}}{e^{V_1} - 1} \right)^{n/2} \cdot \frac{D(\mathcal{R}(y))}{D(y)}$$

where $D : A \to (0, \infty)$ *is given by* $D(y) = |f_0(y) - y|$, n *is the length of the cycle (always even),* V_0, V_1 *are the distortion constants of* f_0, f_1 *respectively.*

Define by $\mathcal{R}_j = h^{-1}_{i_1 \dots i_j}$, $1 \leq j \leq n$, and then $\mathcal{R}_n = \mathcal{R}$. We adopt the convention $\mathcal{R}_0 = \mathrm{id}$. To obtain Proposition 3.6 we need the following two-by-two term estimate.

Lemma 3.7. *For any* $0 \leq j \leq n - 2$ *even,*

$$\frac{|\mathcal{R}_{j+2}(J)|}{|\mathcal{R}_j(J)|} \geq \frac{e^{-(V_0+V_1)}}{e^{V_1} - 1} \cdot \frac{D(\mathcal{R}_{j+2}(y))}{D(\mathcal{R}_j(y))}.$$

This lemma concludes immediately Proposition 3.6. Indeed, we have that

$$\frac{|\mathcal{R}(J)|}{|J|} = \frac{|\mathcal{R}_n(J)|}{|\mathcal{R}_{n-2}(J)|} \cdot \frac{|\mathcal{R}_{n-2}(J)|}{|\mathcal{R}_{n-4}(J)|} \cdots \frac{|\mathcal{R}_2(J)|}{|J|}$$

and applying Lemma 3.7 to each of the terms one gets the estimate required in Proposition 3.6.

Proof of Lemma 3.7. Since

$$\frac{|\mathcal{R}_{j+2}(J)|}{|\mathcal{R}_j(J)|} = \frac{|\mathcal{R}_{j+2}(J)|}{|\mathcal{R}_{j+1}(J)|} \cdot \frac{|\mathcal{R}_{j+1}(J)|}{|\mathcal{R}_j(J)|},$$

the lemma will be proved by estimating each term.

Observe that since the cycle begins with the map f_0, necessarily $j + 2 \pmod 2 = 0$ and $j + 1 \pmod 2 = 1$. Since s_j is a fixed point of f_0 and $\mathcal{R}_j(J)$ is contained in a fundamental domain of f_0 (see Addendum 3.5) then

$$s_j < f_0(\mathcal{R}_j(y)) < \mathcal{R}_j(x) < \mathcal{R}_j(y).$$

Applying $f_0^{-m_{i_1 \dots i_{j+1}}}$ to the fundamental domain $[f_0(\mathcal{R}_j(y)), \mathcal{R}_j(y)]$, the usual distortion estimate gives

$$\frac{|f_0(\mathcal{R}_j(y)) - \mathcal{R}_j(y)|}{|\mathcal{R}_j(x) - \mathcal{R}_j(y)|} \geq e^{-V_0} \frac{|f_0^{-m_{i_1 \dots i_{j+1}}}(f_0(\mathcal{R}_j(y))) - f_0^{-m_{i_1 \dots i_{j+1}}}(\mathcal{R}_j(y))|}{|f_0^{-m_{i_1 \dots i_{j+1}}}(\mathcal{R}_j(y)) - f_0^{-m_{i_1 \dots i_{j+1}}}(\mathcal{R}_j(x))|}.$$

Observe that

$$f_0^{-m_{i_1 \dots i_{j+1}}+1}(\mathcal{R}_j(y)) < s_{j+1} < f_0^{-m_{i_1 \dots i_{j+1}}}(\mathcal{R}_j(y)) \quad \text{and} \quad \mathcal{R}_{j+1} = f_0^{-m_{i_1 \dots i_{j+1}}} \circ \mathcal{R}_j.$$

Using this information and rearranging the terms one obtains,

$$\frac{|\mathcal{R}_{j+1}(J)|}{|\mathcal{R}_j(J)|} \geq e^{-V_0} \frac{|s_{j+1} - \mathcal{R}_{j+1}(y)|}{D(\mathcal{R}_j(y))}.$$

To estimate the other term, since s_{j+2} is again a fixed point of f_0 and $\mathcal{R}_{j+2}(J)$ is contained in a fundamental domain of f_0, it follows

$$s_{j+2} < f_0(\mathcal{R}_{j+2}(y)) < \mathcal{R}_{j+2}(x) < \mathcal{R}_{j+2}(y).$$

By construction of the return map (see Addendum 3.5), $[s_{j+2}, \mathcal{R}_{j+2}(y)] \subset A_{j+2}$, which is a fundamental domain of f_1.

This time let us apply $f_1^{m_{i_1\dots i_{j+2}}}$ to $[f_0(\mathcal{R}_{j+2}(y)), \mathcal{R}_{j+2}(y)]$, contained in a fundamental domain of f_1. Then the distortion estimate becomes

$$\frac{|\mathcal{R}_{j+2}(x)-\mathcal{R}_{j+2}(y)|}{|f_0(\mathcal{R}_{j+2}(y))-\mathcal{R}_{j+2}(y)|} \geq e^{-V_1}\frac{|f_1^{m_{i_1\dots i_{j+2}}}(\mathcal{R}_{j+2}(y))-f_1^{m_{i_1\dots i_{j+2}}}(\mathcal{R}_{j+2}(x))|}{|f_1^{m_{i_1\dots i_{j+2}}}(f_0(\mathcal{R}_{j+2}(y)))-f_1^{m_{i_1\dots i_{j+2}}}(\mathcal{R}_{j+2}(y))|}.$$

Since A_{j+2} is a fundamental domain of f_1,

$$f_1(\mathcal{R}_{j+2}(y)) < f_0(\mathcal{R}_{j+2}(y) < \mathcal{R}_{j+2}(y),$$

then

$$\begin{aligned}|f_1^{m_{i_1\dots i_{j+2}}}(f_0(\mathcal{R}_{j+2}(y)))-f_1^{m_{i_1\dots i_{j+2}}}(\mathcal{R}_{j+2}(y))| <\\ < |f_1^{m_{i_1\dots i_{j+2}}}(f_1(\mathcal{R}_{j+2}(y)))-f_1^{m_{i_1\dots i_{j+2}}}(\mathcal{R}_{j+2}(y))|\end{aligned}$$

As $f_1^{m_{i_1\dots i_{j+2}}} \circ \mathcal{R}_{j+2} = \mathcal{R}_{j+1}$, using the above inequality and rearranging the terms,

$$\frac{|\mathcal{R}_{j+2}(J)|}{|\mathcal{R}_{j+1}(J)|} \geq e^{-V_1}\frac{D(\mathcal{R}_{j+2}(y))}{|f_1(\mathcal{R}_{j+1}(y))-\mathcal{R}_{j+1}(y))|}.$$

And so,

$$\frac{|\mathcal{R}_{j+2}(J)|}{|\mathcal{R}_{j}(J)|} \geq e^{-(V_0+V_1)}\cdot\frac{D(\mathcal{R}_{j+2}(y))}{D(\mathcal{R}_{j}(y))}\cdot\frac{|s_{j+1}-\mathcal{R}_{j+1}(y)|}{|f_1(\mathcal{R}_{j+1}(y))-\mathcal{R}_{j+1}(y))|}.$$

Finally let us estimate the last term. Applying the mean-value theorem to the function $f_1-\mathrm{id}$ and using that s_{j+1} is a fixed point of f_1,

$$\frac{|s_{j+1}-\mathcal{R}_{j+1}(y)|}{|f_1(\mathcal{R}_{j+1}(y))-\mathcal{R}_{j+1}(y))|} = \frac{1}{|Df_1(z)-1|} > \frac{1}{e^{V_1}-1},$$

which proves the lemma. □

Assume the maps f_0, f_1 are close enough to rotations in the $C^{1+\mathrm{bv}}$-topology so that

$$\frac{e^{-(V_0+V_1)}}{e^{V_1}-1} > 1.$$

Notice that, if the distortion constants $V_0, V_1 \leq \varepsilon$, the above condition is fulfilled for every positive $\varepsilon \leq 0.38$. Then also $\left((e^{V_1}-1)e^{(V_0+V_1)}\right)^{-n/2} > 1$. By the same proof of Lemma 2.4, we conclude the following analogous result.

Lemma 3.8. *There exists $N \in \mathbb{N}$ such that $D\mathcal{R}^N(x) > \lambda > 1$ for all $x \in A$.*

3.4. **Proof of Theorem A.** Let us continue assuming that the maps f_i both have fixed points. We will need the next two lemmas.

Lemma 3.9. *There exists a ss-interval for* IFS(Φ) *if and only if there exists a uu-interval.*

Proof. By contradiction, suppose there exists a uu-interval but no ss-intervals (the other case is similar). Then, Lemma 3.3 implies that there is a cycle of length n with the order $s_0 < s_1 < \cdots < s_0 + 1 = s_n$ on the interval $[s_0,\, s_0 + 1]$ identified with the circle.

Consider the maps f_i as maps of the real line (in particular of the interval $[s_0,\, s_0+1]$), and the uu-interval as the interval $[a,\, b] \subset [s_0,\, s_0+1]$. By the definition of the uu-interval, if $x \leq a$, then $f_i(x) \leq a$ and if $x \geq b$, then $f_i(x) \geq b$. And so if $x > b$, then any finite composition of the maps f_0 and f_1, denoted by h, will satisfy that $h(x) > b$.

Take $x > s_{n-1}$ and $x > b$, using the cycle we can inductively attract x to s_{n-2}, then to s_{n-3}, until s_0, such that in the end will obtain a map h, a finite composition of f_0 and f_1, with $h(x) < a$. This contradicts the previous observation. □

Lemma 3.10. *Suppose that there are no ss-intervals for* IFS(Φ) *and for some interval I, $f_i(I) \cap I \neq \emptyset$, $i = 0, 1$. Then for every $x \in S^1$, there exists $h_1, h_2 \in$* IFS(Φ) *such that $h_1(x) \in I$ and $h_2^{-1}(x) \in I$.*

Proof. Since there are no ss-intervals, Lemma 3.3 implies that there is a cycle with the order $s_0 < s_1 < \cdots < s_n + 1$ in the interval $[s_0,\, s_0+1]$. We identify the interval I on the circle S^1 with an interval in $[s_0,\, s_0+1]$ that has the endpoints $a < b$.

Assume first that on the interval $[s_0,\, s_0+1]$, the point x satisfies $b < x \leq s_0+1$. Considering f_i as maps of the interval and arguing as in the proof of the above lemma, by using the attractors of the cycle we can inductively start attracting x to $s_0 = s_n$. Then, there exists h a finite composition of f_0 and f_1 of the form $h = f_i \circ h_1$ ($i = 0$ or 1), such that $h(x) \geq a$ and $h(x) < a$. Since $f_i(I) \cap I \neq \emptyset$, necessarily $h_1(x) \in I$, as required.

For the case when $x \leq a$, suppose $I \subset [s_{i-1}, s_i]$. By means of the cycle but now looking on the circle S^1, we can attract x so s_i so that on the interval $[s_0,\, s_0+1]$, $x > s_i$. Then this is similar to the situation above, applied not to x but to $h(x)$, with $h(x) > s_i \geq b$.

Since by Lemma 3.9 no ss-intervals is equivalent to having no uu-intervals, the same argument can be repeated with IFS(Φ^{-1}) to prove the existence of h_2^{-1} with $h_2^{-1}(x) \in I$. □

3.4.1. *Proof of minimality.* Let us show that IFS(Φ) is minimal. The minimality of IFS(Φ^{-1}) follows since by Lemma 3.9, IFS(Φ^{-1}) has no ss-intervals and then the proof is the same.

Consider a point $x \in S^1$ and an open interval J. Since there is no ss-intervals, by Lemma 3.8, there exists an expanding return map $\mathcal{R}^N$ defined in the domain A and associated with a cycle. Since $A = (s_0,\, f^{-1}(s_0)]$, a fundamental domain of f_1 and is in the basin of attraction of s_0, then $f_i(A) \cap A \neq \emptyset$. By Lemma 3.10, there exists $h_1 \in$ IFS(Φ) with $h_1^{-1}(J) \cap A \neq \emptyset$. We can assume J is small enough so that $h_1^{-1}(J)$ is contained in one of the domains of the map $\mathcal{R}^N$. Since $D\mathcal{R}^N > \lambda > 1$, then $|\mathcal{R}^{kN}(J)| > \lambda^k |J|$. Iterating sufficiently many times $\mathcal{R}^{kN}(J)$ eventually must contain a discontinuity s of $\mathcal{R}^N$.

By Lemma 3.4, $s \in \mathrm{Orb}^+_\Phi(s_0) \cup \cdots \cup \mathrm{Orb}^+_\Phi(s_n)$. Without loss of generality we assume that s is in the orbit of s_0. Then there exists $h_2 \in \mathrm{IFS}(\Phi)$ with $h_2(s_0) = s$ and so

$$h_2(s_0) \in \mathcal{R}^{kN} \circ h_1^{-1}(J).$$

Writing $\mathcal{R}^{kN}$ as h_3^{-1} with $h_3 \in \mathrm{IFS}(\Phi)$, we get $h_1 \circ h_3 \circ h_2(s_0) \in J$.

For the cycle there is the order $s_0 < s_1 < \cdots < s_0 + 1 = s_n$, and thus given a point x we may attract x arbitrary close to s_0 by inductively attracting to each s_i of the cycle. Then there exists h_4 with $h_4(x)$ close enough to s_0. It can be concluded that $h_4 \circ h_1 \circ h_3 \circ h_2(x) \in J$, and since x and J were arbitrary, this proves minimality of $\mathrm{IFS}(\Phi)$.

3.4.2. *Density of hyperbolic periodic points.* Given an interval J, by the previous argument we may suppose there exists $h_1 \in \mathrm{IFS}(\Phi)$ and a point of the cycle, say s_0, such that $h_1(s_0) \in J$. Again by the same reasoning as above, there exists $h_2 \in \mathrm{IFS}(\Phi)$ such that $h_2(J)$ is attracted arbitrary close to s_0. Then we may assume that $h_2 \circ h_1(J) \subset J$ and $D(h_2 \circ h_1)|_J < \beta < 1$. This implies that there is a hyperbolic attractor in J.

The density of hyperbolic repellers for $\mathrm{IFS}(\Phi)$ follows from the density of hyperbolic attractors for $\mathrm{IFS}(\Phi^{-1})$.

3.4.3. *The case of generators with periodic points.* When the generators f_i have non-zero rotation number, consider the next lemma whose demonstration is left to the reader.

Lemma 3.11. *Let f be an orientation preserving $C^{1+\mathrm{bv}}$-diffeomorphism of the circle with periodic points of period n, and consider B the immediate basin of attraction of some periodic point. Then*

$$e^{-V_f} \le \frac{Df^n(x)}{Df^n(y)} \le e^{V_f} \quad \textit{for all } x, y \in B$$

where V_f is the distortion of f on S^1. That is, $\mathrm{Dist}(f^n, B) \le V_f$.

Since in the immediate basin there is always a point with derivative one,

$$e^{-V_f} \le Df^n(x) \le e^{V_f} \quad \textit{for all } x \in S^1.$$

The lemma states that the the distortion of $f_i^{n_i}$ is the same of f_i in the relevant regions and the derivative of of $f_i^{n_i}$ is bounded by the distortion and thus will still be close to one if f_i is close to rotations. Namely, $|Df_i^n(x) - 1| \le e^{V_i} - 1$ where V_i is the distortion constant of f_i. Therefore, the same proof as in the case of fixed points applies. Observe that also in Lemma 3.6 we have to ask again that

$$\frac{e^{-(V_0+V_1)}}{e^{V_1} - 1} > 1.$$

Since $V_0, V_1 \le \varepsilon$, the above condition is fulfilled for all $\varepsilon \le 0.38$.

This completes the proof of Theorem A.

4. Spectral decomposition

4.1. **Spectral decomposition on the real line.** The next theorem gives a complete description of the global topological dynamics of an IFS generated by a pair of diffeomorphisms on the real line.

Theorem 4.1 (Spectral decomposition on the real line)**.** *There exists* $\varepsilon > 0.38$ *such that if* f_0 *and* f_1 *are diffeomorphisms of the real line, one of them having at least a fixed point, with no fixed points in common and* ε*-close to the identity in the* $C^{1+\mathrm{bv}}$*-topology, then*

$$L(\mathrm{IFS}(\Phi)) \setminus \{\pm\infty\} = \bigcup K_i \quad \textit{and} \quad K_i \subset \overline{\mathrm{Per}(\mathrm{IFS}(\Phi))}$$

where each K_i *is a compact, maximal transitive set for* IFS(Φ)*. Moreover, each compact set* K_i *is either*

- *a single fixed point of* f_0 *or* f_1 *or*
- *a* $**$*-interval for* IFS(Φ) *with* $** \in \{ss, su, uu, s, u\}$*.*

In particular if f_0 *and* f_1 *have only hyperbolic fixed points, then the above union of the sets* K_i *is countable and they are pairwise disjoint.*

Proof. Consider $z \in L(\mathrm{IFS}(\Phi)) \setminus \{\pm\infty\}$. We can assume that $z \in \omega(\mathrm{IFS}(\Phi))$ since the situation for the α-limit of IFS(Φ) is followed by a similar argument. Then, by definition of ω-limit of IFS(Φ), the point z is approximated by points of the form $y_k \in \omega_\Phi(x_k)$. For each k, there is a sequence $(h_{kn})_n \subset \mathrm{IFS}(\Phi)$ such that each point y_k is again approximated by points of the form $h_{kn} \circ \dots \circ h_{k1}(x_k)$.

Claim 4.2. *If* $y \in \omega_\Phi(x)$*, then either* y *belongs to some* $**$*-interval for* IFS(Φ) *with* $** \in \{ss, su, uu, s, u\}$ *or it is a fixed point of* f_0 *or* f_1*.*

This claim concludes that the limit set of IFS(Φ) is contained in the union of $**$-intervals and fixed points of f_0 and f_1. Indeed, from Claim 4.2, either, z is a fixed point of f_0 or f_1, or then, for k_0 large enough, y_k belongs in the same $**$-interval for all $k \geq k_0$ and thus z is also in this $**$-interval.

Proof of the claim 4.2. Since $y \in \omega_\Phi(x)$, there is $(h_n)_n \subset \mathrm{IFS}(\Phi)$ such that $y = \lim_{n\to\infty} h_n \circ \dots \circ h_1(x)$. If y is an accumulation point of fixed points then, obviously, y is a fixed point of f_0 or f_1. Thus, we can assume that y is not of this type. Also, without loss of generality we suppose that $y \geq 0$.

Since at least one of the generators has a fixed point, we can assume by the following argument that $y \in [p,\, q]$ where $p, q \in \mathrm{Fix}(f_0) \cup \mathrm{Fix}(f_1)$ and $(p,\, q) \cap \mathrm{Fix}(f_i) = \emptyset$ for $i = 0, 1$. Otherwise, there is a fixed point P such that any other fixed point is less then P and $y \in [P,\, \infty)$. If $y \neq P$, then it is not hard to check via the geometry of the functions, that it is not possible for $f_0, f_1 > \mathrm{id}$ in $(P\, \infty]$, since then $y = \infty$. In the other case, $[P,\, \infty)$ is s or u-interval or $y = P$.

Suppose p and q are both attractors or repellers but for different maps restricted to $[p,\,q]$. As f_0 and f_1 are close to the identity, it follows that this closed interval is a ss or uu-interval.

So, we can consider the final case when p and q are an attractor-repeller pair for the same map, say f_0 (the repeller-atractor case is analogous). Note that then $f_0 < \mathrm{id}$ in $(p,\,q)$. We have two options: $f_1 < \mathrm{id}$ or $f_1 > \mathrm{id}$ in $[p,\,q]$. In the first case, both maps are below the identity and hence this geometry implies that the unique possible ω-limit points for IFS(Φ) in $[p,\,q]$ are p and q. For the second case, we have again two options:

$$f_1([p,\,q]) \cap [p,\,q] \neq \emptyset \quad \text{or} \quad f_1([p,\,q]) \cap [p,\,q] = \emptyset.$$

In the first option $[p,\,q]$ is a su-interval for IFS(Φ), and for the other one it follows as before that y is either p or q.

Therefore we have showed that in every possible case, the point $y \in \omega_\Phi(x)$ belongs to a $**$-interval for IFS(Φ) with $** \in \{ss, su, uu, s, u\}$ or it is a fixed point of f_0 or f_1. □

To conclude the proof of the theorem we need to show every set K_i is contained in the limit set of IFS(Φ). Recall that K_i denotes both, a $**$-interval or a fixed point of f_0 or f_1. It is clear that every fixed point is contained in the limit set and so we only need to prove that every $**$-interval for IFS(Φ) is recurrent.

We will show that there exists $\varepsilon > 0$ such that if f_0 and f_1 are ε-close to the identity in the $C^{1+\mathrm{bv}}$-topology, then any $**$-interval for IFS(Φ) with $** \in \{ss, su, s\}$ satisfies the hypothesis of Theorem 2.2. This proximity implies that $|f_i(x) - x| < \varepsilon$, $|Df_i(x) - 1| < \varepsilon$ and $\mathrm{Dist}(f_i, \mathbb{R}) < \varepsilon$ for $i = 0, 1$, and we only need to demonstrate that $(1-\varepsilon)\varepsilon^{-1}e^{-V} > 1$.

Notice that $V \leq 2\varepsilon|a - f_i^{-1}f_j(a)|$ where $i \neq j$ and the a is the endpoint of the $**$-interval. Suppose a is the left endpoint and $f_0 < \mathrm{id}$ in $(a,\, f_0^{-1}f_1(a)]$. From the proximity to the identity, $f_1(a) \leq a + \varepsilon$. Since

$$f_0(x) \geq (1-\varepsilon)(x-a) + a \quad \text{for all } x \in (a,\, f_0^{-1}f_1(a)],$$

it follows that $f_0^{-1}f_1(a) - a \leq \varepsilon(1-\varepsilon)^{-1}$. Thus,

$$(1-\varepsilon)\varepsilon^{-1}e^{-V} \geq (1-\varepsilon)\varepsilon^{-1}e^{-2\varepsilon^2(1-\varepsilon)^{-1}} > 1$$

for all $\varepsilon \leq 0.38$.

Finally by Theorem 2.2, ss, su and s-intervals are minimal with a dense set of periodic points. In particular, this kind of intervals are transitive sets for IFS(Φ). The same properties are obtained for uu-intervals and u-intervals for IFS(Φ^{-1}).

Lemma 4.3. *If A is minimal for* IFS(Φ^{-1}) *then A is transitive for* IFS(Φ).

This lemma follows from the facts that minimality implies transitivity and if A is transitive for IFS(Φ^{-1}) then A is transitive for IFS(Φ).

Applying the above lemma we obtain that uu and u-intervals are also transitive sets for the IFS(Φ). This basically concludes the proof of the theorem. The last part is immediately obtained since if all the fixed points are hyperbolic, then they are isolated and so there exists only a countable number of the fixed poins and the $**$-intervals. □

Remark 4.4. Theorem 4.1 holds also for IFS generated by a pair of diffeomorphisms on compact interval I under the same assumptions. To see this, we understand the compact interval I like the compactified real line $[-\infty, \infty]$ and the K^s or K^u intervals in I are identified with the relevant intervals on the real line.

4.2. **Spectral decomposition on the circle: Proof of Theorem B.** The following result is the a version of Theorem 2.2 for diffeomorphisms of the circle with periodic points.

Proposition 4.5. *There is $\varepsilon > 0.30$ such that if f_0 and f_1 are circle diffeomorphisms with periodic points of period n_0 and n_1 respectively, ε-close to the rotation in the $C^{1+\mathrm{bv}}$-topology then for every $**$-interval K^{**}, $** \in \{ss, su\}$, with respect to* $\mathrm{IFS}(\Phi^n)$ *it holds that*

$$K^{**} \subset \overline{\mathrm{Per}(\mathrm{IFS}(\Phi^n))} \quad \textit{and} \quad K^{**} \subset \overline{\mathrm{Orb}^+_{\Phi^n}(x)} \quad \textit{for all } x \in K^{**}.$$

Proof. We will apply Theorem 2.2 for $\mathrm{IFS}(\Phi^n)$ for the $**$-interval $K^{**} = [a, b]$. Without loss of generality, we assume that $f_0^{n_0}(x) < x$ for all $x \in (a, b)$. According to Lemma 3.11,

$$e^{-V_0} \leq Df_0^{n_0}(x) \leq e^{V_0} \quad \text{for all } x \in S^1$$

where V_0 is the distortion constant of f_0. This implies that $|Df_0^{n_0}(x) - 1| < e^{V_0} - 1$ for all $x \in S^1$. Similarly we denote by V_1 the total variation of f_1 and then it also follows that

$$\mathrm{Dist}(f_0^{n_0}, I) \leq V_0 \quad \text{and} \quad \mathrm{Dist}(f_1^{n_1}, J) \leq V_1$$

for all fundamental domains I of $f_0^{n_0}$ and J of $f_1^{n_1}$ in $(a, f_0^{-n_0} f_1^{n_1}(a)]$.

Since $(a, f_0^{-n_0} f_1^{n_1}(a)] \subset S^1$, its length is less that one. Let us remember the necessary condition in Theorem 2.2,

$$(1 - (e^{V_0} - 1))(e^{V_0} - 1)^{-1} e^{-(V_0 + V_1)|a - f_0^{-n_0} f_1^{n_1}(a)|} > 1.$$

As $V_0, V_1 \leq \varepsilon$, it suffices to take $\varepsilon \leq 0.30$ to guarantee that the above condition is fulfilled, which concludes the proof of the proposition. □

Proof of Theorem B. This result is immediately achieved from Theorem 4.1, Proposition 4.5 and Theorem A. Indeed, consider $\tilde{f}_0^{n_0}$ and $\tilde{f}_1^{n_1}$ as the lifts on the real line of $f_0^{n_1}$ and $f_1^{n_1}$. Note that $\tilde{f}_0^{n_0}$ and $\tilde{f}_1^{n_1}$ are periodic functions. Arguing as in Theorem 4.1, there is a decomposition in compacts sets on the real line of $L(\mathrm{IFS}(\tilde{f}_0^{n_0}, \tilde{f}_1^{n_1})) \setminus \{\pm\infty\}$. From the periodicity of $\tilde{f}_i^{n_i}$, it follows that each of these compact sets project on the circle as either, a periodic point of f_i, or as a $**$-intervals for $** \in \{ss, su, uu\}$. By Theorem 4.5 these intervals are transitive sets for $\mathrm{IFS}(\Phi^n)$.

What is left is to study the limit set of point whose ω-limit (or α-limit) contains $\pm\infty$ on the real line. This can only happen if there is a cycle for $\mathrm{IFS}(\Phi^n)$ different of a ss-interval. Then Theorem A implies that S^1 is minimal for $\mathrm{IFS}(\Phi^n)$ and $\mathrm{IFS}(\Phi^{-n})$, a contradiction. Therefore, we obtain a decomposition of the limit set, as required. □

5. A Denjoy type Theorem for IFS

5.1. **A trichotomy for IFS.**

The next lemma shows some relations between the ω-limit sets and the orbits of an IFS. These properties will be necessary afterwards for the proof of the trichotomy result. We will use the following notation: given a set A on a manifold M, we denote by A' the set of accumulation points of A, i.e., the set of points y such that there exists a sequence $(x_n)_n \subset A$ converging to y with $x_n \neq y$ for all $n \in \mathbb{N}$.

Lemma 5.1. *Consider $\Phi = (\phi_1, \ldots, \phi_k) \in \mathrm{Hom}(M)^k$, a non-empty closed subset K of a manifold M. Then it holds that:*

i) $\omega_\Phi(h(x)) \subset \omega_\Phi(x) \subset \overline{\mathrm{Orb}^+_\Phi(x)}$ for all $x \in M$ and for $h \in \mathrm{IFS}(\Phi)$,

ii) if $K = \overline{\mathrm{Orb}^+_\Phi(x)}$ for all $x \in K$ then

$$K = \bigcup_{i=1}^{k} \phi_i(K) = \omega_\Phi(x) \quad \textit{for all } x \in K,$$

iii) $\mathrm{Orb}^+_\Phi(x)' = \phi_1(\mathrm{Orb}^+_\Phi(x)') \cup \cdots \cup \phi_k(\mathrm{Orb}^+_\Phi(x)')$ for all $x \in M$.

Proof. By definition of the ω-limit set of a point $x \in M$, $\omega_\Phi(h(x)) \subset \omega_\Phi(x)$ for all $h \in \mathrm{IFS}(\Phi)$. On the other hand, since

$$\overline{\mathrm{Orb}^+_\Phi(x)} = \{y : \text{there exists } (g_n)_n \subset \mathrm{IFS}(\Phi) \text{ such that } y = \lim_{n\to\infty} g_n(x)\}$$

then $\omega_\Phi(x)$ is a subset of the closure of the orbit of x. Therefore, we conclude (i).

According to the first item, to obtain (ii) it suffices to prove that

$$K \subset \bigcup_{i=1}^{k} \phi_i(K) \quad \text{and} \quad K \subset \omega_\Phi(x) \quad \text{for all } x \in K.$$

The first inclusion is followed from the minimality of K. Indeed, for any $y \in K$ there is $(g_n)_n \subset \mathrm{IFS}(\Phi)$ such that $x = \lim_{n\to\infty} g_n(y)$. Since we are working with a finitely generated semigroup, taking a subsequence if necessary, we can assume that $g_n = \phi_i \circ \tilde{g}_n$ for some $i \in \{1, \ldots, k\}$ and $\tilde{g}_n \subset \mathrm{IFS}(\Phi)$. Thus, $\phi_i^{-1}(x) \in K$ as it is the limit of $\tilde{g}_n(y)$ and K is an invariant closed set. Therefore $x = \phi_i(\phi_i^{-1}(x)) \in \phi_i(K)$, showing the first inclusion.

In order to prove the second inclusion, we fix $x, y \in K$ and consider a sequence of positive real numbers $\varepsilon_n = 1/n \to 0$. Let us construct by induction a sequence $(h_n)_n \subset \mathrm{IFS}(\Phi)$, such that the distance between y and $h_n \circ \cdots \circ h_1(x)$ is less than ε_n. Since the orbit of x is dense in K, there is $h_1 \in \mathrm{IFS}(\Phi)$ such that $d(y, h_1(x)) < \varepsilon_1$. Similarly, since $h_1(x) \in \mathrm{Orb}^+_\Phi(x) \subset K$ then the orbit of $h_1(x)$ is dense in K and there exists h_2 such that $d(y, h_2 \circ h_1(x)) < \varepsilon_2$. Inductively we obtain the desired sequence $(h_n)_n \subset \mathrm{IFS}(\Phi)$. Hence, $y = \lim_{n\to\infty} h_n \circ \cdots \circ h_1(x)$ and thus $y \in \omega_\Phi(x)$ for all $x, y \in K$, concluding (ii).

Now we will show that $\mathrm{Orb}^+_\Phi(x)'$ is a self-similar set, that is prove the last item. Note that $\phi_i(\mathrm{Orb}^+_\Phi(x)') \subset \mathrm{Orb}^+_\Phi(x)'$ for all $i = 1, \ldots, k$. Indeed, if y is an accumulation point of the orbit of x, then $\phi_i(y)$ is approximated by points of the form $\phi_i \circ g_n(x) \in \mathrm{Orb}^+_\Phi(x)$

where $y = \lim_{n\to\infty} g_n(x)$ and $g_n(x) \neq y$ for all $n \in \mathbb{N}$. This implies that $\phi_i(y)$ is also an accumulation point of $\mathrm{Orb}^+_\Phi(x)$ and so we conclude one of the inclusions.

To obtain the other inclusion

$$\mathrm{Orb}^+_\Phi(x)' \subset \phi_1(\mathrm{Orb}^+_\Phi(x)') \cup \cdots \cup \phi_k(\mathrm{Orb}^+_\Phi(x)'),$$

we fix any $y \in \mathrm{Orb}_\Phi(x)'$ and then there exists $(g_n) \subset \mathrm{IFS}(\Phi)$ such that $y = \lim_{n\to\infty} g_n(x)$ and $g_n(x) \neq y$ for all $n \in \mathbb{N}$. Since the semigroup $\mathrm{IFS}(\Phi)$ is finitely generated, taking a subsequence if necessary, we can assume that for some fixed $i \in \{1,\dots,k\}$ we have $g_n = \phi_i \circ \tilde{g}_n$ with $\tilde{g}_n \in \mathrm{IFS}(\Phi)$. Hence, $\phi_i^{-1}(y) = \lim_{n\to\infty} \tilde{g}_n(x)$ and $\tilde{g}_n(y) \neq \phi_i^{-1}(y)$. Since the accumulation set is a closed set, it holds that $\phi_i^{-1}(y) \in \mathrm{Orb}^+_\Phi(x)'$. This implies that $y \in \phi_i(\mathrm{Orb}^+_\Phi(x)')$ proving the desired inclusion and therefore (iii). □

The next trichotomy theorem is the IFS counterpart of the similar result for group actions in [5]. An interval I is forward-invariant for $\mathrm{IFS}(\Phi)$ if $f_i(I) \subset I$ for all I.

Theorem 5.2. *(Trichotomy for IFS) Consider* $\mathrm{IFS}(\Phi)$ *generated by* $\Phi = (f_1,\dots,f_k) \in \mathrm{Hom}(I)^k$, *where* I *is* S^1 *or a forward-invariant compact interval. Then there exists a non-empty closed subset* K *of* M *such that*

$$K = \bigcup_{i=1}^{k} f_i(K) \quad \text{and} \quad K = \overline{\mathrm{Orb}^+_\Phi(x)} = w_\Phi(x) \ \text{for all } x \in K.$$

Moreover, one (and only one) of the following possibilities occurs:

- *i) K is a finite orbit for* $\mathrm{IFS}(\Phi)$,
- *ii) K has non-empty interior,*
- *iii) K is a Cantor set.*

Proof. A family of non-empty closed subsets of M such that for each member Λ, $\Lambda = f_1(\Lambda) \cup \cdots \cup f_N(\Lambda)$ can be ordered by inclusion. Since the intersection of nested compact sets is compact and non-empty, Zorn's Lemma allows to conclude the existence of a minimal (regarding the inclusion) non-empty closed set K such that $K = \cup f_i(K)$. We will show that K is an invariant minimal set for $\mathrm{IFS}(\Phi)$,

$$K = \overline{\mathrm{Orb}^+_\Phi(x)} \quad \text{for all } x \in K. \tag{5}$$

Then, according to Item (ii) in Lemma 5.1, we have $K = \overline{\mathrm{Orb}^+_\Phi(x)} = \omega_\Phi(x)$ for all $x \in K$.

In order to prove (5), since K is an closed invariant set, $\mathrm{Orb}^+_\Phi(x)' \subset \overline{\mathrm{Orb}^+_\Phi(x)} \subset K$ for all $x \in K$. On other hand, according to (iii) in Lemma 5.1, for each $x \in K$ the set of accumulation points $\mathrm{Orb}^+_\Phi(x)'$ is a closed self-similar set. Since K is minimal (regarding the inclusion) then either, $\mathrm{Orb}^+_\Phi(x)'$ is an empty set or $K = \mathrm{Orb}^+_\Phi(x)'$. We have two possibilities:

(i) there is $x \in K$ such that $\mathrm{Orb}^+_\Phi(x)'$ is empty;
(ii) for all $x \in K$, it holds that $K = \mathrm{Orb}^+_\Phi(x)'$.

In the first case, it follows that $\mathrm{Orb}^+_\Phi(x)$ is finite set, and therefore it is a non-empty closed self-similar set contained in K. This implies, via Zorn's Lemma, that $K = \mathrm{Orb}^+_\Phi(x) = \omega_\Phi(x)$.

In the second case, we obtain that K is an invariant minimal set for IFS(Φ) and so $K' = K$. Moreover, we have two options: K has non-empty interior or the interior of K is empty and thus, since M is a one-dimensional manifold, K is a Cantor set. This concludes the proof of the theorem. □

As an attractor for IFS is generated by a forward-invariant set, the theorem above describes the shape of possible attractors of an IFS.

Remark 5.3. In the case of I is a ss-interval, which is forward-invariant, the end-points are global (with respect to I) attractors and are necessarily part of any minimal set in I. Thus the minimal set is unique. See also [14].

5.2. **Proof of Theorem C.** From Theorem B it follows that any closed invariant minimal set, Λ, for IFS(Φ^n) is contained in some piece of the spectral decomposition of the limit set. By the hypothesis of the theorem this piece cannot be a single point because the generators do not have fixed points in common. It also cannot be a su or a uu-interval because they are not forward invariant and any point can be made to leave the interval. Thus necessarily the piece has to be a ss-interval K^{ss}, in which case by Remark 5.3 , the only invariant minimal set is K^{ss} and so $\Lambda = K^{ss}$.

It is left to show that there cannot exist invariant Cantor sets for IFS(Φ). Let us suppose that Λ is an invariant minimal Cantor set for IFS(Φ) and take any point $p \in \Lambda$. Since f_0 and f_1 have no periodic points in common, the basins of all attracting periodic points of f_0 and f_1 cover S^1. By hypothesis S^1 is not minimal and so from Theorem A there is at least one ss-interval for IFS(Φ^n) whose endpoints are attractors. The cover of S^1 by the basins of attractors allows p to be attracted inside some ss-interval K^{ss} for IFS(Φ^n). Since Λ is an invariant minimal set for IFS(Φ), then $K^{ss} \subset \Lambda$, contradicting that it has empty interior.

5.3. **A Theorem of Duminy on group actions of the circle.** Finally, we will show Duminy's Theorem [13] under the assumption of Theorem A. We denote by G(Φ^n) the group generated by $\Phi^n = (f_0^{n_0}, f_1^{n_1})$.

Theorem 5.4 (Duminy)**.** *There exists $\varepsilon > 0.30$ such that if f_0 and f_1 are diffeomorphisms on the circle with periodic points of period n_0 and n_1 respectively, ε-close to the rotations in the $C^{1+\mathrm{bv}}$-topology and with no periodic points in common then,*

- *S^1 is minimal for G(Φ^n) and*
- *the hyperbolic periodic points of G(Φ^n) are dense in S^1.*

Proof. Since there are no periodic points in common and we are working with a group then considering the inverses if necessary, there always exists a ss-interval and a cycle different of the ss-interval for the group G(Φ^n). Moreover, this cycle has the same order as in Lemma 3.3. By Proposition 4.5 every point in the ss-interval has a dense orbit for the action of G(Φ^n). Now similarly to the proof of minimality in Theorem B, the cycle allows to bring any point and interval to the ss-interval and this implies minimality of S^1 for G(Φ^n).

The density of hyperbolic periodic points follows from the argument described in Lemma 2.5 (see also Section 3.4.2). This concludes the proof of the result. □

ACKNOWLEDGEMENTS

Both authors thank Enrique R. Pujals for useful conversations and P. G. Barrientos is especially grateful to José A. Rodríguez for the orientation and the helpful comments.

During the preparation of this article the authors were partially supported by the following fellowships: P. G. Barrientos by FPU-grant AP2007- 031035, MTM2008-06065 project (Spain) and CNPq post-doctoral fellowship (Brazil); A. Raibekas by CNPq doctoral fellowship (Brazil) and CAPES-PNPD (Brazil) pos-doctoral fellowship. We thank Instituto Nacional de Matemtica Pura e Aplicada (IMPA) for its hospitality.

REFERENCES

[1] P. G. Barrientos, Y. Ki, and A. Raibekas. Symbolic blender-horseshoes and applications. *ArXiv e-prints*, Nov. 2012.

[2] C. Bonatti, L. J. Díaz, and M. Viana. *Dynamics beyond uniform hyperbolicity*, volume 102 of *Encyclopedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2005.

[3] A. Denjoy. Sur les courbes définies par les équations différentielles à la surface du tore. *J. de Math Pures Apll.*, 11:333–375, 1932.

[4] G. Duminy. L'invariant de godbillon-vey d'un feuilletage se localise dans les feuilles ressort. Université de Lille, 1970.

[5] É. Ghys. Groups acting on the circle. *Enseign. Math. (2)*, 47(3-4):329–407, 2001.

[6] A. S. Gorodetskiĭ and Y. S. Il′yashenko. Some new robust properties of invariant sets and attractors of dynamical systems. *Funktsional. Anal. i Prilozhen.*, 33(2):16–30, 95, 1999.

[7] A. S. Gorodetskiĭ and Y. S. Il′yashenko. Some properties of skew products over a horseshoe and a solenoid. *Tr. Mat. Inst. Steklova*, 231(Din. Sist., Avtom. i Beskon. Gruppy):96–118, 2000.

[8] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.

[9] V. Kleptsyn and D. Volk. Skew products and random walks on intervals. *ArXiv e-prints*, Oct. 2011.

[10] V. A. Kleptsyn and M. B. Nal′skiĭ. Convergence of orbits in random dynamical systems on a circle. *Funktsional. Anal. i Prilozhen.*, 38(4):36–54, 95–96, 2004.

[11] M. Nassiri and E. R. Pujals. Robust transitivity in hamiltonian dynamics. *Ann. Sci. Éc. Norm. Supér.*, 45(2), 2012.

[12] A. Navas. Sur les groupes de difféomorphismes du cercle engendrés par des éléments proches des rotations. *Enseign. Math. (2)*, 50(1-2):29–68, 2004.

[13] A. Navas. *Groups of circle diffeomorphisms*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, spanish edition, 2011.

[14] K. Shinohara. On minimality of semigroup actions on interval which are C^1-close to the identity. *ArXiv e-prints*.

[15] M. Viana and J. Yang. Physical measures and absolute continuity for one-dimensional center direction. *Annales de l'Institut Henri Poincare (C) Non Linear Analysis*, to appear.

Departamento de Matemática PUC-Rio
Marquês de So Vicente 225, Gávea, Rio de Janeiro 225453-900, Brazil

E-mail address: `barrientos.pg@gmail.com`

Instituto de Matemática e Estatística, UFF
Rua Mário Santos Braga s/n - Campus Valonguinhos, Niterói, Brazil

E-mail address: `artem@impa.br`